

Mapping coastal aquaculture and fisheries structures by satellite imaging radar

Case study of the Lingayen Gulf, the Philippines

by

Carlo Travaglia

Environment and Natural Resources Service
FAO Sustainable Development Department
Rome, Italy

Giuliana Profeti

SAR Image Processing
Florence, Italy

José Aguilar-Manjarrez

Inland Water Resources and Aquaculture Service
FAO Fisheries Department
Rome, Italy

and

Nelson A. Lopez

Bureau of Fisheries and Aquatic Resources
Manila, the Philippines

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

ISBN 92-5-105114-3

All rights reserved. Reproduction and dissemination of material in this information product for educational or other non-commercial purposes are authorized without any prior written permission from the copyright holders provided the source is fully acknowledged. Reproduction of material in this information product for resale or other commercial purposes is prohibited without written permission of the copyright holders. Applications for such permission should be addressed to:

Chief

Publishing Management Service

Information Division

FAO

Viale delle Terme di Caracalla, 00100 Rome, Italy

or by e-mail to:

copyright@fao.org

© FAO 2004

FOREWORD

Cooperation between the FAO Remote Sensing Center (now Environment and Natural Resources Service (SDRN)) and the Inland Water Resources and Aquaculture Service (FIRI) began in 1983 with the joint planning of the international training course “Remote Sensing Applications to Inland Fisheries and Aquaculture”, held at FAO headquarters in September 1984. Its aim was to raise awareness among fishery technical and management officers of the very positive remote sensing and GIS applications which were relevant to their work. In this framework and with the same objectives in the subsequent years training courses, pilot studies and operational projects have been jointly conducted worldwide.

A result of this longstanding cooperation has been the development and field testing, under operational conditions, of new remote sensing methodologies for specific fisheries requirements, such as aquaculture site selection, wetlands monitoring, shrimp farms inventory and monitoring and others. These methodologies have been widely disseminated to potential users in both the fields of fisheries and remote sensing through FAO publications.

The paper we present here aims to introduce a cost-effective new methodology for accurate inventory and monitoring of coastal aquaculture and fisheries structures.

Jiansan Jia

Chief, Inland Water Resources and
Aquaculture Service (FIRI)

Jeff Tschirley

Chief, Environment and Natural Resources
Service (SDRN)

Distribution:

FAO Members and Associate Members
FAO Fisheries Department
FAO Fisheries Officers in FAO Regional and Subregional Offices
FI Marine Fisheries and Aquaculture Mailing Lists

ACKNOWLEDGEMENTS

The authors are greatly indebted to all who assisted in the implementation and completion of this study by providing information, advice and facilities.

The administrative support of the Bureau of Fisheries and Aquatic Resources (BFAR), the Philippines, Director Malcolm I. Sarmiento, Jr. and of the other Bureau's officers at the regional and provincial level is hereby acknowledged. Likewise the field assistance of Leodegario Abarra, Nestor Botacion, Catherine Rabadon, Mea Baldonado, Lina Bravo, Felita Ugaban, Roberto De Vera, Jr. and Wilfredo Basa is gratefully appreciated, as well as the participation in the field team of the BFAR-IFAD Technical Staff; James Villanueva, Rodolfo Tigue, Roy Garcia and Edgardo Amurao.

The provision by the European Space Agency (ESA) of satellite radar data at nominal cost, acquired according to the authors specifications, greatly facilitated the work and has been sincerely appreciated.

The authors are greatly indebted to Roberto Carlà of the Institute of Applied Physics of the Italian National Research Council; Pierre-Philippe Mathieu, Earth Observation Science and Applications, European Space Agency; James McDaid Kapetsky, Senior Fishery Resources Officer (FAO-FIRI retired) and Uwe Barg, Fishery Resources Officer (FAO-FIRI) for their critical review of the manuscript.

Claudia Riva assisted in preparing data for the field verification work and in the literature review. Desktop publishing work was by Nadia Pellicciotta.

Travaglia, C.; Profeti, G.;
Aguilar-Manjarrez, J.; Lopez, N.A.
Mapping coastal aquaculture and fisheries structures by satellite imaging radar.
Case study of the Lingayen Gulf,
the Philippines.
FAO Fisheries Technical Paper. No. 459. Rome, FAO. 2004. 45p.

ABSTRACT

Inventry and monitoring of coastal aquaculture and fisheries structures provide important baseline data for decision-making in planning and development, including regulatory laws, environmental protection and revenue collection. Mapping these structures can be performed with good accuracy and at regular intervals by satellite remote sensing, which allows observation of vast areas, often of difficult accessibility, at a fraction of the cost of traditional surveys.

Satellite imaging radar (SAR) data are unique for this task not only for their inherent all-weather capabilities, very important as aquaculture activities mainly occur in tropical and subtropical areas, but essentially because the backscatter from the structure components allows for their identification and separation from other features.

The area selected and object of the study has been Lingayen Gulf, sited in Northwestern Luzon Island, the Philippines, where all these structures of interest occur.

Field verification of the methodology resulted in the following accuracy: fishponds 95 percent, fish pens 100 percent. Mapping accuracy for fish cages was estimated at 90 percent and for fish traps at 70 percent.

The study is based on interpretation of SAR satellite data and a detailed image analysis procedure is described. The report aims at the necessary technology transfer for an operational use of the approach indicated in other similar environments.

Keywords: Aquaculture; Fisheries structures; Geographic Information Systems; Lingayen Gulf; Philippines; Remote Sensing; SAR; Satellite imaging radar.

CONTENTS

	Page
Foreword	iii
Acknowledgements	iv
Abstract	v
List of Figures	viii
List of Tables	ix
1. Introduction	1
1.1 Background: the Sri Lanka experience on mapping inland aquaculture farms	1
1.2 Objective of the present study	3
1.3 Description of the test area	3
1.4 Description of the structures: fish pens, cages and traps	5
2. Methodology	9
2.1 Data and software used	9
2.2 Preparation of the vector database	11
2.3 Mapping aquaculture and fisheries structures mapping by satellite imaging radar	12
2.4 Image pre-processing procedure	19
2.5 Digitalization of the shoreline	19
2.6 Mapping procedures	20
3. Results	23
3.1 Fishponds	24
3.2 Fish pens	25
3.3 Fish cages	26
3.4 Fish traps	26
3.5 Land cover changes	28
3.6 Field verification exercise	28
4. Discussion	33
4.1 Radar fine beam	33
4.2 ERS SAR	33
4.3 Final considerations and recommendations	33
5. Glossary	35
References	39
Recommended further reading	41
Appendix	44
Map of coastal aquaculture and fisheries structures in Lingayen Gulf, the Philippines	45

LIST OF FIGURES

	Page
1. Shrimp farms in the Dutch Canal test site, Sri Lanka	2
2. The study area and the zones covered by the satellite data	4
3. Fish pens in Lingayen Gulf	6
4. Fish cages in Lingayen Gulf	7
5. Fish traps in Lingayen Gulf	8
6. Topographic maps, 1:50 000 scale, covering the study area	10
7. A portion of Dagupan City topographic map (sheet 7074 II) and the derived vector data	13
8. Interaction of radar beams with dykes and water surfaces on a group of fishponds	14
9. Return signal as a function of the angle between a dyke and the cross-track direction	15
10. The angle between the scanning directions in the SAR data used in the study	16
11. Interactions of radar beams with a fish cage	17
12. Sea state and coastal aquaculture and fisheries structures mapping	18
13. Interpreted RADARSAT-1 SAR image and the resulting map of the aquaculture and fisheries structures	24
14. Colour combination of the three SAR images used in the study	25
15. Appearance on RADARSAT-1 and ERS-2 images of small metallic and non-metallic fish cages	26
16. Appearance of large fish cages on RADARSAT-1 SAR data	27
17. Appearance of offshore fish traps on the RADARSAT-1 SAR image	27
18. Appearance of fish traps inside rivers on the RADARSAT-1 SAR image	28
19. Position of the verification points and of some observation points	30

LIST OF TABLES

1.	Total surface covered by shrimp farms, Sri Lanka study	2
2.	Coordinates of the study area	3
3.	Characteristics of the satellite data used	9
4a.	Classes identified in the SAR images: Polygon layer	20
4b.	Classes identified in the SAR images: Polyline layer	21
5.	Total area covered by classes of interest (Pangasinan province)	23
6.	Length of the fish traps detected in the study area	23
7.	Comparison between the fishponds area coverage in 1977 and in 2002	25
8.	Comparison of visual interpretation and <i>in situ</i> observations on the verification points	31

